



Entrant's Report

Harvest 2022

YEN Field ID: BF22EXAMPLE

Entrant name: EXAMPLE

Main contact email:
EXAMPLE

Sponsor/supporter: EXAMPLE

Sponsor/Supporter email:
EXAMPLE

Field/Site name: EXAMPLE







Incident energy: NA TJ/ha

Available water: NA mm

Crop: Winter bean

Variety: EXAMPLE

SUMMARY: YEN entries were completed from 8 winter and 25 spring crops out of 39 original entries, this year we are classing crops by their sowing season rather than variety type. Headline results for your entry are shown in benchmark diagrams below. Your yield of NA t/ha represents NA% of its estimated yield potential of NA t/ha.



CONTENTS

Our detailed analysis of your yield result is provided in the following pages, including comparisons with other YEN entries, and with benchmarks derived from previous YEN data where possible. We hope that this helps you to identify aspects of your husbandry and growing conditions that offer possible routes to further yield enhancement.

Our approach in this report is to consider growing conditions and potential yields for crops grown in this season, then the conditions for and husbandry of your crop, its development, its basic resources (light energy, water & nutrients), its success in capturing these and in converting them to seed. Lastly, we use seed analysis to provide a post-mortem on your crop's limiting components and nutrition.

The benchmarking diagrams in this report only include the data set submitted by the YEN data submission deadline. Reports produced using data submitted after this deadline show an entrants value in comparison to this previously reference data set.

CONTENTS	2
POTENTIAL SEED YIELDS	3
2022 Potential yields	3
SEASONAL GROWING CONDITIONS	4
Soil description and nutrition analysis	5
Soil analysis	6
AGRONOMY	7
Husbandry factors	7
Husbandry factors continued	9
CROP DEVELOPMENT	10
Spring	10
Winter	11
RESOURCES AND THEIR CAPTURE	12
Water availability and capture	12
Energy capture	13
Nutrient capture	14
YIELD ANALYSIS	16
Yield formation	16
Yield components	18
Seed formation and size	19
CROP NUTRITION POST-MORTEM	20
SUMMARY	22
The 2022 Bean YEN competition:	22
UPCOMING EVENTS	22
CONTACTS	22
YEN SPONSORS	22

POTENTIAL SEED YIELDS

"The YEN exists to help you to enhance your yields."


To estimate potential yields, we assume a theoretically 'perfect' variety grown with 'inspired' husbandry on your land with this season's weather, achieving either:

- (i) **70% capture of light energy** through this season (including some in September), and its conversion to 1.2 tonnes of biomass per terajoule, or
- (ii) **Capture of all the available water** held in the soil to 1 m depth (or to rock if shallower than 1 m) plus all rainfall from April to July, and conversion of each 25 mm into a tonne of biomass per hectare. Our model of potential yield estimates potential growth on a daily basis; this identifies impacts of water limitation more precisely than the cruder monthly estimates we made in previous YEN reports.

Taking the lesser of these two biomass amounts, we assume that a maximum of 60% can be used to form seed, this is the 'harvest index'. Note that we assume average temperatures for the UK, and no damage from waterlogging, frost, heat, or lodging.

The maps below show the potential seed yields for autumn sown cereals on retentive and light soils this year. For this we assume deep soils with no irrigation.

2022 Potential yields



We are using weather data from DTN™ this year. Note we do not have long term met data from DTN so cannot show a map of long-term average yield potentials.

SEASONAL GROWING CONDITIONS




The adjacent graphs show the monthly temperatures, rainfall and total solar radiation for your area through this growing season compared to your regional long-term average (LTA) and the average for all UK arable areas (1981-2010, from the Met Office).

The winter was milder than average, and whilst January was dry February was very wet. Winter beans were mostly drilled successfully, with generally mild temperatures allowing timely drilling. The spring was also warmer than average, with March warmer and brighter than usual. The relatively dry March and April allowed Spring bean growers to drill crops into moisture, but the following dry spell may have affected many crops.

The summer was warmer than average and maximum temperatures were far above the usual for most of England. There was little rain for most of July and early August, although localized thunderstorms contributed to rainfall in some areas. The summer of 2022 was also sunnier than usual.


The lack of rain and high temperatures meant that crops were at least two weeks earlier in the South of the country compared to previous years. Some winter beans were ready by the start of August and Spring beans in the middle of August. The lack of rain depressed yields especially in spring beans but also kept diseases to a minimum.

Stem nematode was much less prevalent this season due to the dry weather. Bruchid infestation on the other hand was much higher, mainly due to the average temperature rarely being below 20 °C. The warmer temperatures also suited black bean aphids.



YEN Benchmarking charts – What do they mean?

YEN provides a full set of metrics whereby you can gauge the performance of your crop against all other YEN crops. This has provided the principle value of YEN to most participants. We do this with benchmark charts. These compare your value with everyone else's this year and with standard benchmarks and critical values, if available and appropriate. The key is as follows:



The 'whiskers' show the range of YEN values in 2022 whilst the grey box shows the middle half of values, with a line for the mid-value. The orange line shows the value for your entry, and the red line is a limit beyond which yield may be adversely affected; crops with values beyond this merit further investigation.

Note that 'Dynamic Benchmarking' is available to all YEN members via the YEN website. This means you can compare your own yield or seed nutrient data with subsets of all other YEN crops selected by crop type, soil type, location or year back to 2013.

Soil description and nutrition analysis




Your soil's capacity to hold available water is critical in determining your potential yields. We rely on entrants describing the soil where their YEN entry grew. We can use the [UK Soil Observatory map viewer](#) to check whether this complies with the surrounding land.


Good soil descriptions are vital in allowing us to estimate soil water holding capacity and, along with summer rainfall, the water available to your crop (see Benchmark charts in the section on 'Resources & their Capture').

Topsoil analyses provided by NRM also tell us about soil status for pH, P, K and Mg, as reported on the next page. A few sites show low values for soil pH, P, K or Mg. If these are unexpected, they may need further checks, either by repeating soil analysis and by checking both leaf and seed analyses later in this report. Previous YEN leaf and seed nutrient data have indicated that UK cereal crops often experience deficiencies in one or more nutrients, and sometimes this is despite soil levels being satisfactory. So, by combined use of soil, leaf and seed analysis, the YENs now help to diagnose whether nutrient shortfalls are arising from poor supply, or poor capture by the root system.


Soil OM, %




Soil pH




Soil P, mg/l




Soil K, mg/l




Soil Mg, mg/l




Sand, % w/w



Silt, % w/w



Clay, % w/w



SOM supports crop performance through better nutrient availability, soil aggregation, and water holding capacity. NRM determines SOM by 'loss on ignition'. Note: other methods can give lower values.

Soil pH <6 is acid. High pH soils may require that special attention is paid to phosphorus (P) and micro-nutrient levels in the leaf and seed (see later).

Only a small difference separates P Index 0 (≤ 9) and 2 (≥ 16). High yields are possible at P index 1 but fresh P is also usually required. Use seed P (see page 20) to double-check if P was sufficient.

Soil potassium (K) analysis checks on whether K supplies are likely to have been deficient for average crops. However, high yielding crops require very large amounts of K.

Magnesium (Mg) is a key component of chlorophyll so deficient plants show striking inter-veinal yellowing. Temporary deficiencies often occur in spring if topsoils are dry.

Soils with high sand content hold the least water and soils with high silt content tend to hold the most water.


Soils with much silt and sand, hence less clay, tend to be relatively weak, and so are more difficult to maintain with a stable structure.

Soils with high clay content hold much water but part of this is held too tightly for crop use. Nutrients within this unavailable water tend to be less available than nutrients in lighter soils.

AGRONOMY

This section considers how your variety and husbandry decisions related to others entering the YEN this year. The most chosen varieties are compared in the figure below for their maturity and seed protein levels, as reported in the PGRO's Descriptive list.

- Your variety was EXAMPLE.




Husbandry factors


The following diagrams use orange segments or orange bars to indicate the agronomy of your crop, if known, so you can see how this relates to all other YEN entries. Analysis of all Bean YEN entries from 2019-2021 shows the following associations with seed yield (note that these do not necessarily imply causes – it may just be that farms with high yields also happen to have these traits):

Soil type	Better yields on soils with lower clay content. As winters in the Bean YEN have generally been wet, this may suggest potential difficulties of establishment in some heavy soils and improvements to soil structure could be beneficial to yields
Soil analysis	Better yields with more soil P
K inputs and seed K	Better yields with more K
Fungicide use	Higher yields associated with good disease control
Harvest losses	Higher harvest losses associated with lower yields


Main cultivation strategy




Previous Crop Type




Predominant organic materials applied




Area wildflowers mixes on farm




Type of woodlands/hedgerows nearby



Establishment of nearby woodland/hedgerow habitats




Proportion of flowering crops in 1 mile radius




Husbandry factors continued


Sowing date, winter varieties




Number of herbicides applied




Sowing date, spring varieties




Seeds sown per m²




Number of insecticides applied



Number of fungicides applied




Fertiliser P₂O₅ applied, kg/ha




Fungicide spend, £/ha




Fertiliser K₂O applied, kg/ha




Crop protection spend, £/ha



Fertiliser SO₃ applied, kg/ha



Crop protection spend, £/tonne



CROP DEVELOPMENT


The following charts show how your entry developed through the 2021-22 season, compared to all other Bean YEN entries and Benchmarks. The cardinal stages of emergence (GS10), flowering (GS61) and full senescence (GS89) determine the length of each phase for growth:

- Foundation, GS10-GS31 – when development of leaves and side shoots occurs;
- Construction, GS31-GS61 – when stem elongation occurs, and flowers are formed;
- Production, GS65-GS89 – when pods are formed, and seeds are filled.

We have separated out the dates of development for spring and winter sown crops over the next two pages.


Spring

Emergence date, spring




Crops are sorted into winter and spring by sowing date rather than variety type.


Nodulation (GS34)




Flowering (GS61)




Canopy senescence (GS87)



Harvest date



Crop height, cm



These are the average heights from 10 shoots in each grab sample.


Height of the lowest pod, cm



These are the average heights from 10 shoots in each grab sample.


Winter

Emergence date, winter




Crops are sorted into winter and spring by sowing date rather than variety type.


Nodulation (GS34)




Flowering (GS61)




Canopy senescence (GS87)



Harvest date




Crop height, cm



These are the average heights from 10 shoots in each grab sample.

Height of the lowest pod, cm




These are the average heights from 10 shoots in each grab sample.

RESOURCES AND THEIR CAPTURE

Water availability and capture


This page shows how weather this year affected the water available for your crop and other crops entered in the YEN. Water is supplied through the main growing period from concurrent rainfall and also from water stored in the soil. UK soils almost always refill with water over-winter. Water potentially available to each crop through the summer includes this soil water plus the summer rainfall (April to July).

Soil water holding capacity, mm



Deep soils hold water to a great depth; we assume roots can access all easily held water (to 2 bar suction) to a depth of 1 m (or to rock, if shallower). If enough roots didn't reach to this depth, capture of soil-available water will have been accordingly less.


Rainfall April–July, mm



After winter drainage stops, spring and summer rainfall is held in the topsoil until it is evaporated or transpired by the crop's canopy.


Whilst we cannot yet measure water captured by YEN crops individually, by assuming your crop's conversion of water to total biomass was 'normal' (25 mm water for each t/ha biomass formed), we have made crude estimates below of the likely success of your crop's root system in capturing water.

Total water available, mm



Total water is the sum of your soil's water-holding capacity and your summer rainfall (both shown above).

Estimated use of available water, %



Small water use will sometimes have been due to less demand for canopy transpiration (e.g. because crop developed faster and matured earlier) or otherwise due to worse rooting.

If your estimated use of available water exceeds the total water available, this may be good news! It either suggests that your crop's roots were more efficient than normal, or that your soil description was overly pessimistic: i.e. your soil apparently managed to provide more water than we estimated was possible from your soil's texture, stone content and depth.


A high yielding crop, growing say 9.4 t/ha of biomass (so yielding 5 t/ha seed at 45% harvest index and 15% moisture), would need to capture ~240 mm water from soil reserves plus summer rain.

Energy capture

The benchmarking charts below show how weather this year affected light energy available for this entry and other YEN crops. Solar radiation has been divided into periods that roughly equate to the three key phases of crop development reported above:


- Foundation – when development of leaves and branches occurs (Winter beans October to March; Spring beans March to May (fewer side shoots formed))

Solar radiation Oct–Mar, TJ/ha




- Construction – when stem elongation occurs, and flowers are formed (Winter beans March to May; Spring beans April to June)

Solar radiation Apr–May, TJ/ha




- Production – when pods are formed, and seeds are filled (Winter beans May to September; Spring beans June to October)

Solar radiation Jun–July, TJ/ha




Whilst we cannot yet measure light capture by YEN crops individually, by assuming your crop's conversion of light-energy was 'normal' (1 tonnes/TJ), we have made a crude estimate below of the likely success of your crop's canopy in capturing total light-energy for the 12 months of this season.

Solar radiation total, TJ/ha/yr



Total solar radiation across YEN entries is generally less in the north and more in the south.

Estimated % solar radiation captured





Average light capture tends to be poor if a crop's lifespan is short. The biophysical limit for beans is possibly two thirds.

Nutrient capture

Whether nutrient capture was sufficient to support full conversion of light and water is best deduced from nutrient concentrations in crop tissues – both leaves (next three pages) and seeds (later section). No critical thresholds or benchmarks are shown for leaf analyses because these change through a crop's life and are still uncertain. However, the benchmarking diagrams should enable you to compare your crop's levels with all other YEN entries this year, analysed at the same time.

Lancrop Laboratories provide leaf analyses for YEN. Samples are of the newest fully expanded leaf.





YIELD ANALYSIS


Yield formation

The whole-crop samples that YEN entrants provide all have their components counted and weighed and results are shown in the following charts, assuming that each sample was representative of the whole area from which seed yield was determined. [Area-related values such as seeds/m² are derived from the validated yield are derived from the validated seed yield.]

Total biomass production indicates the success with which a crop captured its key resources, light-energy and water, and the harvest index (the proportion of total biomass that was harvestable) indicates how this biomass was apportioned to seed. Since seed growth happens last, harvest index also indicates how late growth related to early growth.


Your seed yield (expressed as t/ha and % of potential) is shown below along with biomass and harvest index, in relation to all other YEN entries.

Above-ground biomass, t/ha




The average winter bean crop biomass in 2022 was 8.9 t/ha. The average spring bean crop biomass in 2022 was 6.7 t/ha. Experience with other crops shows that high biomass relates to high yields.

Harvest index, %




The average harvest index for winter bean crops in 2022 was 52%. The average value for spring crops was 57%.

Seed yield, t/ha




The average value for winter crops was 5.5 t/ha. The average value for spring crops was 4.5 t/ha

Straw Yield, t/ha




Straw Yields here include stubble and pod walls. The average value for winter crops was 4.3 t/ha. The average value for spring crops was 2.9 t/ha.

Estimated harvest loss, t/ha




Estimated from the number of seeds seen on the ground after harvest, and the average weight per seed.

Seed yield potential, t/ha



YEN yield potential express the light energy and water available for your entry this year, simply converted to t/ha. The average value for winter crops was 8.7 t/ha. The average value for spring crops was 8.6 t/ha.

% yield potential



Bean YEN yields in 2022 averaged at 62% of yield potential for winter crops and 53% for spring crops.


Yield components

Whole crop yield analysis can also tell us about the history of your crop because the different crop components are determined sequentially, at least within given layers of the canopy for beans. So comparing your yield components with those of other YEN entrants should indicate the stage(s) through the season at which your crop deviated from others and from normal.

Total biomass provides the overall summary of your crops' growth; this depends on plant numbers determined first, shoot numbers determined next, pod numbers determined next, seed numbers determined next and the seed size determined last.


Analysis of 2019-2021 Bean YEN data suggest that high yields are associated with high pods/shoot, seeds/pods and seeds/m² rather than shoots per plant, however, high yields can be built in different ways.

Plants No./m²




Plants per m² are calculated from the plant counts or overhead photos you provided. If you did not provide plant counts some calculated metrics below might be missing. The average was 36 and 28 for winter and spring crops respectively.

Shoots per plant




Shoots per plant were counted from your grab sample. Note that if plants were cut above ground level this could reduce the value. The average was 1.66 and 1.05 for winter and spring crops respectively.

Pods per shoot




Pod numbers were counted on 10 shoots. Taller stems may have space for more pods per shoot. This count may have included infertile pods. The average was 12 and 14 for winter and spring crops respectively.

Pods /m²




Pods per m² is a calculation of pods per shoot and shoots per plant. The average was 271 and 326 for winter and spring crops respectively.

Seeds per pod



The expected value is 3-5 seeds per pod. Infertile pods in the grab sample may lead to underestimation. The average was 3.1 and 2.6 for winter and spring crops respectively.

Seeds /m²



Seeds per m² in the crop are a reflection of seed set. Here, seeds per m² are estimated from you combine seed sample, TSW and crop yield. The average was 715 and 717 for winter and spring crops respectively.


Seed formation and size

We use your combine-harvested seed sample to provide the analysis of seed size and seed filling. Seed filling depends mainly on photosynthesis after flowering, therefore it largely relies on the health and longevity of the green canopy.


If the number of seeds per m^2 is low (see above), or if conditions during early seed-fill are limiting, final seed filling, hence yield may be constrained even if later conditions are good – this is sometimes described as ‘sink’ limitation. Sink limitation is often more important for beans than for wheat.

Bruchid beetle can affect both winter and spring bean varieties. Adults emerge from seeds leaving a circular hole, staining and larvae in the seeds. Beetles do not breed in seed stores, but damaged beans may not be accepted for quality markets. Bruchid damage can also cause abnormal seedlings and increased infection by soil-borne pathogens such as damping off, so reducing germination and maybe also the vigour of the seedlings.


Combine TSW, g (15%MC)



Seed protein, %



Bruchid damage, %




Bruchid damage affects acceptance for quality markets, and maybe also germination and seedling vigour.



CROP NUTRITION POST-MORTEM

The YEN has trail-blazed use of seed analysis to provide overall post-mortems on a crop's nutrition.

- Results from >900 YEN cereal samples analysed up to last year suggest that nutrient deficiencies are very common (using the 8 critical values that we know so far); >80% of crops showed deficiencies, and >50% showed two or more deficiencies! Phosphorus deficiency has been most common.
- YEN Nutrition was therefore launched in 2021 to help to remedy these deficiencies – further details and registration are available [here](#)
- Crop nutrients differ in how they are shared between grain and straw at harvest. The graph below shows how different crop species store most of their N & P in the seed but most of their K in the straw (as estimated from analyses of feed materials).



- We now use YEN-low values (i.e. lower quartiles from all past YEN data – the boundary between the bottom quarter and top three-quarters of all YEN values since 2013) as comparators (red lines) for all nutrients in all crops. We find YEN-low values to be very similar to known critical thresholds of N, P, S and Mn in wheat, as well as to less certain critical values of K, Mg, Cu & Zn, so we assume they can be applied for all nutrients in all crops.
- The following benchmarking-charts and YEN-low values provide the best means of identifying the nutrient(s) most likely to have limited your crop.



SUMMARY

The 2022 Bean YEN competition:

- Thank you for providing the samples and information necessary to complete this report; the collective efforts of all YEN contributors maximise the value of the results that can be reported and the deductions that can be made.
- The highest seed yield in 2022 was 7.5 t/ha (in Scotland). This winter bean crop also achieved the highest percentage of biophysical potential at 83%.

UPCOMING EVENTS

‘YEN is ten’ Conference If you haven't done so already, please register and join us to celebrate the YEN's tenth birthday at a cross-YEN conference at East of England Arena, Peterborough on 24th January 2023. You can Register for the YEN Conference [here](#).

Pea and Bean YEN 2022 discussion meeting

In addition to the cross crop YEN Conference 2022 we are pleased to invite Pea YEN and Bean YEN members and sponsors to a Pulse YEN specific meeting to discuss the season and learnings from the Pulse YEN data sets on the 16th January 2022 at PGRO, The Research Station, Great North Road, Thornhaugh, Peterborough PE8 6HJ. More details and a link to register are to follow. We hope to see you there!

CONTACTS

Please send any comments, observations or queries to the contacts below.

Thomas Wilkinson	Thomas.Wilkinson@adas.co.uk	07814043347
Charlotte White	Charlotte.White@adas.co.uk	07503570264

Or email yen@adas.co.uk for general enquiries.

[@adasYEN](#)

YEN SPONSORS

The YEN was initiated by industry and is entirely industry funded. We are most grateful to all our sponsors. They not only provide funding but they are fundamentally involved in management of the YEN and in supporting individual farms in making their YEN entries. The YEN would not exist without them!



Visit www.yen.adas.co.uk for sponsors' details, news updates and to register for 2022.